

NZY DNase I

Catalogue number MB19901
Presentation 200 U

Description

NZY DNase I is a highly specific recombinant DNase from bovine pancreas recombinantly produced in *Pichia pastoris*. The enzyme is used for the efficient removal of contaminating DNA. RNA integrity is unchanged after treatment with NZY DNase I under recommended conditions, since it presents no detectable RNase activity.

Shipping & Storage Conditions

Upon arrival, this product can be stored from -15 °C to -25 °C and is stable till the expiry date if stored as specified. After resuspension, store from -15 °C to -25 °C.

Instructions for preparation

NZY DNase I is dissolved in 100 µL of RNase-free water. Incubate for 1 min at room temperature. Gently swirl the vial to completely dissolve the enzyme. NZY DNase I is sensitive to mechanical agitation. Dispense into aliquots and store from -15 °C to -25 °C. The frozen working solution is stable for 6 months. Do not freeze/thaw the aliquots more than three times.

Standard Protocol

In general, the commonly used RNA purification methods co-purify DNA to a considerable extent (e.g., phenol-based RNA purification). This often requires a subsequent removal of contaminating DNA and clean-up of the RNA from the reaction mixture. To remove DNA from such preparations please proceed as follows:

1. Prepare a reaction mixture with 1 µL of enzyme and 10 µL of the 10x Buffer
2. Add a 1/10 volume of the enzyme:buffer reaction to the crude RNA solution (RNA solution should be free of RNase activity).
3. Incubate for 10 min at 37 °C.
4. Repurify the RNA following a suitable clean up procedure.

For on-column digestion

Prepare DNase I working solution as below

	Volume (µl)
DNase I (2 Units/µl)	2
10 x DNase Buffer	5
RNase free water	43

1. Add 50 µl of DNase I working solution on the central of membrane.
2. Incubate at room temperature for 15 min.

Quality Control

Purity

NZY DNase I purity is >95% as judged by SDS polyacrylamide gel electrophoresis followed by Coomassie Blue staining.

Nuclease assays

To test for RNase contamination, 1 µg of RNA is incubated with 1 µL of enzyme for 1 hour at 37 °C. Following incubation, the nucleic acid is visualized on a GreenSafe-stained agarose gel. There must be no visible nicking or cutting of the RNA.

For life science research only. Not for use in diagnostic procedures.